

# Curvature of the Heisenberg Group

Bartek Ewertowski and M. W. Wong

**Abstract.** We compute the Riemannian curvature of the Heisenberg group and then contract it to the sectional curvature, Ricci curvature and the scalar curvature of the Heisenberg group. The main result so obtained is that the Heisenberg group is a space of constant positive scalar curvature.

**Mathematics Subject Classification (2000).** 53C21.

**Keywords.** Heisenberg group, left-invariant vector fields, Riemannian metric, Levi–Civita connection, Riemannian curvature, sectional curvature, Ricci curvature, scalar curvature.

## 1. The Heisenberg Group

If we identify  $\mathbb{R}^2$  with the complex plane  $\mathbb{C}$  via the obvious identification

$$\mathbb{R}^2 \ni (x, y) \leftrightarrow z = x + iy \in \mathbb{C},$$

and we let

$$\mathbb{H}^1 = \mathbb{C} \times \mathbb{R},$$

then  $\mathbb{H}^1$  becomes a noncommutative group when equipped with the multiplication  $\cdot$  given by

$$(z, t) \cdot (w, s) = \left( z + w, t + s + \frac{1}{4}[z, w] \right), \quad (z, t), (w, s) \in \mathbb{H}^1,$$

where  $[z, w]$  is the symplectic form of  $z$  and  $w$  defined by

$$[z, w] = 2 \operatorname{Im}(z\bar{w}).$$

In fact,  $\mathbb{H}^1$  is a unimodular Lie group on which the Haar measure is just the ordinary Lebesgue measure  $dz dt$ .

---

This research has been supported by the Natural Sciences and Engineering Council of Canada under Discovery Grant 0008562.

Let  $\mathfrak{h}$  be the Lie algebra of left-invariant vector fields on  $\mathbb{H}^1$ . Then a basis for  $\mathfrak{h}$  is given by  $X$ ,  $Y$  and  $T$ , where

$$X = \frac{\partial}{\partial x} + \frac{1}{2}y \frac{\partial}{\partial t},$$

$$Y = \frac{\partial}{\partial y} - \frac{1}{2}x \frac{\partial}{\partial t}$$

and

$$T = \frac{\partial}{\partial t}.$$

It can be checked easily that

$$[X, Y] = -T$$

and all other commutators among  $X$ ,  $Y$  and  $T$  are equal to 0.

The aim of this paper is to prove that the scalar curvature of the Heisenberg group is a positive number. This is achieved by contracting from the Riemannian curvature to the scalar curvature through the sectional curvature and the Ricci curvature. The interest in curvature of the Heisenberg group  $\mathbb{H}^1$  stems from the fact [?] that  $\mathbb{H}^1$  can be thought of as the three-dimensional surface that is the boundary of the four-dimensional Siegel domain, so curvature of the Heisenberg group  $\mathbb{H}^1$  may be of some interest in physics.

Results on curvature of the Heisenberg group exist in the literature with a host of different notation and convention. See, for instance, [2, 6, 7]. This paper is another attempt to give a self-contained, elementary and streamlined account based on the notions that can be found in any graduate textbook on Riemannian geometry.

Since indices permeate Riemannian geometry, we find it convenient to label the vector fields  $X$ ,  $Y$  and  $T$  by  $X_1$ ,  $X_2$  and  $X_3$  respectively.

## 2. The Riemannian Metric

We begin with the fact that there exists a left-invariant Riemannian metric  $g$  on  $\mathbb{H}^1$  that turns  $X_1$ ,  $X_2$  and  $X_3$  into an orthonormal basis for  $\mathfrak{h}$  with respect to an inner product denoted by  $(\cdot, \cdot)$ . In fact,

$$g(x, y, t) = \begin{bmatrix} 1 + (y^2/4) & -xy/4 & -y^2/2 \\ -xy/4 & 1 + (x^2/4) & x/2 \\ -y^2/2 & x/2 & 1 \end{bmatrix}$$

for all  $(x, y, t) \in \mathbb{H}^1$ . Equivalently, the arclength variable  $ds$  is given by

$$ds^2 = dx^2 + dy^2 + \left( dt - \frac{ydx - xdy}{2} \right)^2.$$

### 3. The Levi–Civita Connection

A *connection*  $\nabla$  on  $\mathbb{H}^1$  is a mapping

$$\mathfrak{h} \times \mathfrak{h} \ni (X, Y) \mapsto \nabla_X Y \in \mathfrak{h}$$

such that

- $\nabla_X(Y + Z) = \nabla_X Y + \nabla_X Z$ ,
- $\nabla_{X+Y}Z = \nabla_X Z + \nabla_Y Z$ ,
- $\nabla_X(fY) = X(f)Y + f\nabla_X Y$ ,
- $\nabla_{fX}Y = f\nabla_X Y$

for all vector fields  $X, Y$  and  $Z$  in  $\mathfrak{h}$  and all  $C^\infty$  real-valued function  $f$  on  $\mathbb{H}^1$ . The *torsion*  $T$  of the connection  $\nabla$  is a mapping that assigns to two vector fields  $X$  and  $Y$  in  $\mathfrak{h}$  another vector field  $T(X, Y)$  in  $\mathfrak{h}$  given by

$$T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y].$$

A connection  $\nabla$  on  $\mathbb{H}^1$  is said to be *compatible* with the Riemannian metric  $g$  on  $\mathbb{H}^1$  if

$$X(Y, Z) = (\nabla_X Y, Z) + (Y, \nabla_X Z) \quad (3.1)$$

for all vector fields  $X, Y$  and  $Z$  in  $\mathfrak{h}$ .

**Remark 3.1.** In (3.1),  $(Y, Z)$  is a real-valued function, so  $X(Y, Z)$  is also a real-valued function, and the equation expresses an equality of real-valued functions.

The following result is crucial.

**Theorem 3.2.** *There exists a unique affine connection  $\nabla$  on  $\mathbb{H}^1$  such that  $\nabla$  is torsion-free, i.e.,*

$$T(X, Y) = 0$$

for all vector fields  $X$  and  $Y$  in  $\mathfrak{h}$  and  $\nabla$  is compatible with the Riemannian metric  $g$  on  $\mathbb{H}^1$ .

**Proof** The metric  $g$  is given by

$$g_{ij} = (X_i, X_j), \quad i, j = 1, 2, 3.$$

By compatibility,

$$X_i g_{jk} = (\nabla_{X_i} X_j, X_k) + (X_j, \nabla_{X_i} X_k), \quad i, j, k = 1, 2, 3. \quad (3.2)$$

Since  $\nabla$  is torsion-free, it follows that

$$\nabla_{X_i} X_j - \nabla_{X_j} X_i = [X_i, X_j], \quad i, j = 1, 2, 3.$$

Permuting the indices in (3.2), we obtain

$$X_j g_{ik} = (\nabla_{X_j} X_i, X_k) + (X_i, \nabla_{X_j} X_k) \quad (3.3)$$

$$X_k g_{ij} = (\nabla_{X_k} X_i, X_j) + (X_i, \nabla_{X_k} X_j). \quad (3.4)$$

By (3.2), (3.3) and (3.4), we get

$$\begin{aligned} & X_i g_{jk} + X_j g_{ik} - X_k g_{ij} \\ = & 2(\nabla_{X_i} X_j, X_k) - ([X_i, X_j], X_k) + ([X_j, X_k], X_i) - ([X_k, X_i], X_j). \end{aligned} \quad (3.5)$$

Thus, the uniqueness of  $\nabla$  follows. For the existence, For  $i, j, k = 1, 2, 3$ , let

$$\Gamma_{ij}^k = \frac{1}{2} \sum_{l=1}^3 (X_i g_{jl} + X_j g_{il} - X_l g_{ij}) g^{lk},$$

where  $[g^{jk}]$  is the inverse of  $[g_{ij}]$ , and we define  $\nabla_{X_i} X_j$  by

$$\nabla_{X_i} X_j = \sum_{k=1}^3 \Gamma_{ij}^k X_k.$$

□

The connection alluded to in Theorem 3.2 is known as the *Levi-Civita connection*. The functions  $\Gamma_{ij}^k$  are called the Christoffel symbols. We shall work with the Levi-Civita connection from now on.

#### 4. The Riemannian Curvature

Let  $\nabla$  be the Levi-Civita connection on  $\mathbb{H}^1$ . Then the Riemannian curvature  $R$  on  $\mathbb{H}^1$  is the mapping that assigns three vector fields  $X, Y$  and  $Z$  in  $\mathfrak{h}$  another vector field in  $\mathfrak{h}$  denoted by  $R(X, Y)Z$  and given by

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z. \quad (4.1)$$

**Remark 4.1.** An intuitive way to think of the Riemannian curvature  $R$  is that it measures the deviation of  $\nabla_X \nabla_Y - \nabla_Y \nabla_X$  from  $\nabla_{[X, Y]}$ . It should be noted that the opposite sign of  $R$  is also common in the literature. For example, the sign used in [5] is different from the one used in this paper

The Riemannian curvature has many symmetries as given by the following theorem, which can be proved easily using (4.1).

**Theorem 4.2.** *Let  $X, Y, Z$  and  $W$  be in  $\mathfrak{h}$ . Then we have the following symmetries.*

- $R(X, Y)Z + R(Y, X)Z = 0$ ,
- $(R(X, Y)Z, W) + (R(X, Y)W, Z) = 0$ ,
- $R(X, Y)Z + R(Y, Z)X + R(Z, X)Y = 0$ ,
- $(R(X, Y)Z, W) = (R(Z, W)X, Y)$ .

In order to perform computations on the Heisenberg group  $\mathbb{H}^1$ , the following theorem is very useful.

**Theorem 4.3.** *For all vector fields  $X, Y$  and  $Z$  in  $\mathfrak{h}$ , we have*

$$(\nabla_X Y, Z) = \frac{1}{2} \{ (Z, [X, Y]) - (Y, [X, Z]) - (X, [Y, Z]) \}.$$

**Proof** By compatibility,

$$X(Y, Z) = (\nabla_X Y, Z) + (Y, \nabla_X Z) = 0,$$

$$Y(X, Z) = (\nabla_Y X, Z) + (X, \nabla_Y Z) = 0,$$

$$Z(X, Y) = (\nabla_Z X, Y) + (X, \nabla_Z Y) = 0.$$

Since  $\nabla$  is torsion-free, we use the Jacobi identity to the effect that

$$[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$$

to get

$$0 = 2(\nabla_X Y, Z) - (Z, [X, Y]) + (Y, [X, Z]) + (X, [Y, Z]),$$

as asserted.  $\square$

The following two theorems can be proved by direct computations.

**Theorem 4.4.** *The Levi-Civita connection  $\nabla$  is given by*

$$\begin{aligned} \nabla_{X_1} X_1 &= 0, & \nabla_{X_1} X_2 &= -\frac{1}{2} X_3, & \nabla_{X_1} X_3 &= \frac{1}{2} X_2, \\ \nabla_{X_2} X_1 &= \frac{1}{2} X_3, & \nabla_{X_2} X_2 &= 0, & \nabla_{X_2} X_3 &= -\frac{1}{2} X_1, \\ \nabla_{X_3} X_1 &= \frac{1}{2} X_2, & \nabla_{X_3} X_2 &= -\frac{1}{2} X_1, & \nabla_{X_3} X_3 &= 0. \end{aligned}$$

We can now give a useful formula for the Riemannian curvature.

**Theorem 4.5.** *For all vector fields  $X, Y$  and  $Z$  in  $\mathfrak{h}$ ,*

$$\begin{aligned} R(X, Y)Z &= -\frac{3}{4}((Y, Z)X - (X, Z)Y) \\ &\quad + (Y, X_3)(Z, X_3)X - (X, X_3)(Z, X_3)Y \\ &\quad + (X, X_3)(Y, Z)X_3 - (Y, X_3)(X, Z)X_3. \end{aligned}$$

## 5. The Sectional Curvature

Let  $X$  and  $Y$  be two orthonormal vector fields in  $\mathfrak{h}$ . Then  $X$  and  $Y$  determine a *plane* in  $\mathfrak{h}$ . Using left translations, we get a *plane bundle* on  $\mathbb{H}^1$ . Let  $(z, t) \in \mathbb{H}^1$ . Then we can find a neighborhood  $U$  of the origin in  $T_{(z,t)}\mathbb{H}^1$  and a neighborhood of  $(z, t)$  in  $\mathbb{H}^1$  such that the exponential mapping  $\exp : U \rightarrow N$  is a diffeomorphism. As such, the *plane* (a subspace of  $T_{(z,t)}\mathbb{H}^1$ ) induces a submanifold of  $\mathbb{H}^1$  locally and its curvature is given by the so-called *sectional curvature* that we can now define.

**Definition 5.1.** Let  $X$  and  $Y$  be orthonormal vector fields in  $\mathfrak{h}$ . Then the sectional curvature  $S(X, Y)$  determined by  $X$  and  $Y$  is a number given by

$$S(X, Y) = (R(X, Y)X, Y).$$

We can now compute the sectional curvature of the Heisenberg group.

**Theorem 5.2.** *Let  $X$  and  $Y$  be orthonormal vector fields in  $\mathfrak{h}$ . Then*

$$S(X, Y) = \frac{3}{4} - (X, X_3)^2 - (Y, X_3)^2.$$

**Proof** By Theorem 4.5,

$$\begin{aligned} & R(X, Y)X \\ = & -\frac{3}{4}[(Y, X)X - (X, X)Y] \\ & + (Y, X_3)(X, X_3)X - (X, X_3)(X, X_3)Y \\ & + (X, X_3)(Y, X)X_3 - (Y, X_3)(X, X)X_3. \end{aligned}$$

So,

$$(S(X, Y) = (R(X, Y)X, Y) = \frac{3}{4} - (X, X_3)^2 - (Y, X_3)^2.$$

□

## 6. The Ricci Curvature

Let  $X$  and  $Y$  be vector fields in  $\mathfrak{h}$ . Then we consider the linear mapping

$$\mathfrak{h} \ni Z \mapsto R(X, Z)Y \in \mathfrak{h}.$$

We denote this mapping by  $M(X, Y) : \mathfrak{h} \rightarrow \mathfrak{h}$  and we define the *Ricci curvature*  $r(X, Y)$  of the Heisenberg group by

$$r(X, Y) = \text{tr } M(X, Y).$$

**Theorem 6.1.** *Let  $X$  and  $Y$  be vector fields in  $\mathfrak{h}$ . Then the Ricci curvature of  $r(X, Y)$  of the Heisenberg group is given by*

$$r(X, Y) = \frac{1}{2}(X, Y) - (X, X_3)(Y, X_3).$$

**Proof** Using the orthonormal basis  $X_1, X_2$  and  $X_3$  for  $\mathbb{H}^1$ , we get by means of Theorem 4.5

$$\begin{aligned} & R(X, X_j)Y \\ = & -\frac{3}{4}[(X_j, Y)X - (X, Y)X_j] \\ & + (X_j, X_3)(Y, X_3)X - (X, X_3)(Y, X_3)X_j \\ & + (X, X_3)(X_j, Y)X_3 - (X_j, X_3)(X, Y)X_3. \end{aligned}$$

So, for  $j = 1, 2, 3$ ,

$$\begin{aligned}
& (R(X, X_j)Y, X_j) \\
= & -\frac{3}{4}[(X_j, Y)(X, X_j) - (X, Y)] \\
& + (X_j, X_3)(Y, X_3)(X, X_j) - (X, X_3)(Y, X_3) \\
& + (X, X_3)(X_j, Y)(X_3, X_j) - (X_j, X_3)(X, Y)(X_3, X_j).
\end{aligned}$$

Therefore by Parseval's identity,

$$\begin{aligned}
& \text{tr } M(X, Y) \\
= & -\frac{3}{4}[(X, Y) - 3(X, Y)] \\
& + (X, X_3)(Y, X_3) - 3(X, X_3)(Y, X_3) \\
& + (X, X_3)(Y, X_3) - (X, Y) \\
= & \frac{1}{2}(X, Y) - \frac{5}{4}(X, X_3)(Y, X_3),
\end{aligned}$$

as required.  $\square$

By Theorem 6.1, the Ricci curvature is the mapping  $\text{Ric} : \mathfrak{h} \rightarrow \mathfrak{h}$  given by

$$\text{Ric}(X) = \frac{1}{2}X - (X, X_3)X_3$$

for all  $X$  in  $\mathfrak{h}$ .

## 7. The Scalar Curvature

The scalar curvature  $\kappa$  of the Heisenberg group is defined by

$$\kappa = \text{tr}(\text{Ric}).$$

**Theorem 7.1.** *The scalar curvature  $\kappa$  of the Heisenberg group is given by*

$$\kappa = \frac{1}{2}.$$

**Proof** Let  $X_1, X_2$  and  $X_3$  be an orthonormal basis for  $\mathbb{H}^1$ . Then

$$\begin{aligned}
\kappa &= \text{tr}(\text{Ric}) \\
&= \sum_{j=1}^3 (\text{Ric } X_j, X_j) \\
&= \frac{1}{2} \sum_{j=1}^3 (X_j, X_j) - \sum_{j=1}^3 (X_j, X_3)^2 \\
&= \frac{3}{2} - 1 = \frac{1}{2}.
\end{aligned}$$

$\square$

## References

- [1] A. Arvanitoyeorgos, *An Introduction to Lie Groups and the Geometry of Homogeneous Spaces*, American Mathematical Society, 2003.
- [2] L. Capogna, D. Danielli, S.D. Pauls and J. T. Tyson, *An Introduction to the Heisenberg Group and the Sub-Riemannian Isoperimetric Problem*, Birkhäuser, 2007.
- [3] A. Dasgupta, S. Molahajloo and M. W. Wong, The spectrum of the sub-Laplacian on the Heisenberg group, *Tohoku Math. J.* **63** (2011), 269–276.
- [4] A. Dasgupta and M. W. Wong, The semigroup and the inverse of the Laplacian on the Heisenberg group, *CUBO A Mathematical Journal* **12** (2010), 83–97.
- [5] M. P. do Carmo, *Riemannian Geometry*, Birkhäuser, 1992.
- [6] B. Ewertowski, *The Riemannian Geometry of the Heisenberg Group*, M.A. Survey Paper, York University, 2014.
- [7] J. Milnor, Curvatures of left invariant metrics on Lie Groups, *Adv. in Math.* **21** (1976), 293–329.
- [8] G. Segal, Lie Groups, in *Lectures on Lie Groups and Lie Algebras*, Editors: R. Carter, G. Segal and I. Macdonald, London Mathematical Society, 1995, 45–132.
- [9] M. W. Wong, *Partial Differential Equations: Topics in Fourier Analysis*, CRC Press, 2014.

Bartek Ewertowski  
Department of Mathematics  
The University of Auckland  
Private Bag 92019  
Auckland 1142  
New Zealand  
e-mail: bewe542@aucklanduni.ac.nz

M. W. Wong  
Department of Mathematics and Statistics  
York University  
4700 Keele Street  
Toronto, Ontario M3J 1P3  
Canada  
e-mail: mwwong@mathstat.yorku.ca